

Anticancer effect of an olive tree extract through his cytotoxic, antioxidant and antiangiogenic activities

Laure Eloy², Thierry CRESTEIL^{2,3}, Jamal GHANAM¹, Wafa LAABOUDI¹, and Mohammed BENLEMLIH¹

³ IPSIT, Faculté de Pharmacie, Université Paris Sud, 92290 Chatenay-Malabry, France.

² ICSN-CNRS UPR 2301, Avenue de la terrasse, 91190 Gif sur Yvette, France.

¹ Biotechnology Laboratory, Faculty of Science Dhar El Mahraz University Sidi Mohamed Ben Abdellah, P.O. Box 1796 Atlas, Fez-Morocco.

Abstract

The aim of the current study was to evaluate the anticancerous potential of a phenolic olive tree extract through his cytotoxic, antioxidant and antiangiogenic activities. The *in vitro* cytotoxic activity of a crude olive tree extract and his major constitutive components has been evaluated against human KB, HL60 cancer cell lines using MTS and flow cytometry. ROS production was estimated with the DCFH-DA assay. Antiangiogenic activity was evaluated *in vitro* on endothelial cells tube formation and the pro-angiogenic factor expression was quantified using qRT-PCR. *In vitro* cellular assays have demonstrated the cytotoxic effect of the crude olive tree extract. This extract reduces significantly ($p < 0.05$) ROS produced in cells exposed to oxidative stress. Beside this, olive tree extract has demonstrated a strong anti-angiogenic activity, which was correlated with a significant decrease ($p < 0.05$) in VEGF, angiopoietin and HIF1 α expression. Basically, the evaluation of anti-proliferative, antioxidant and anti-angiogenic activities could be the first step to formulate an efficient pharmaceutical product with preventive and/or curative properties against cancer.

Key words: Olive tree extract; Anticancerous; Cytotoxic; Antioxidant; Antiangiogenic.

Introduction

Olive is a generic name for about 35 species of evergreen shrubs and trees of *Olea* genus and *Oleaceae* family, native to tropical and warm temperate regions. The name is particularly used to describe *Olea europaea* L. space cultivated for its edible fruits (olives). The use of olive fruits and leaves for alimentary, therapeutic and cosmetic purposes was a well-known practice in the Mediterranean basin for more than 7 000 years (Evagelia et al., 2004). Although polyphenols are present in olive leaves, olive fruits has been known for several years as a major source of phenolic compounds with high biological activities (Japón-Luján et al., 2007; Mario-Casas et al., 2003; Visioli et al., 2000). Actually, combined of olive fruits and leaves (10% of the total weight of the olives produced generated during olive harvest and olive tree size) to biophenols extracting enhance both qualitative and quantitative extraction yield.

The phenolic fraction of olive tree (fruits, leaves and very young leaves) is a complex mixture of chemical compounds with different structures and molecular weight obtained using mainly liquid-liquid extraction (methanol-water). Olive tree polyphenols belong generally to the following classes: (a) tyrosol, hydroxytyrosol and their secoiridoide derivatives; (b) 4-hydroxybenzoic acid, 4-hydroxy-phenylacetic acid and 4-hydroxycinnamic acid derivatives; (c) lignans and (d) flavonoids. These various compounds have antioxidant (Nardia et al., 2014; Benavente-García et al., 2000), antimicrobial (Micol et al., 2005) and antiproliferative (Taamalli et al., 2012) properties. Today, data collected from *in vitro*, *in vivo* and clinical studies reveal the benefits of these bioavailable compounds for human health. These antioxidant compounds are useful agents for preventing cellular aging and age-related diseases by improving mitochondrial function (Raederstorff et al., 2010). The benefits of olive phenolic compounds on the cardiovascular system have been well documented (Covas, 2007, 2006; Esturch et al., 2006). Beside, these compounds also modulate the inflammatory response associated to several diseases (Killeen et al., 2014; Scoditti et al., 2012; Richard at al., 2011).

Published data have reported that apoptosis and/or necrosis are implicated in the cytotoxicity of olive tree polyphenols. Thus, it has been demonstrated that oleuropein and hydroxytyrosol induce apoptosis in HT-29 human colon adenocarcinoma, HL-60 promyelocytic leukemia) and MCF-7 breast cancer cell lines (Fabiani et al. 2002,

2006, Han et al., 2009). More recently, LeGendre et al. (2015) stated that oleocanthal rapidly and selectively induces apoptotic or necrotic processes via lysosomal membrane permeabilization. Beside this, through an *in vivo* study, Hamdi and Castellon (2005) showed that oleuropein inhibits tumor growth in tumor-bearing mice. Additionally, many epidemiological studies suggest the possible correlation between the consumption of olive products and the incidence of breast cancer (Giacosa et al., 2013).

In the present study, we investigated the cytotoxic effect of an olive tree extract against several cancer cell lines through commonly used cellular assays. Additionally, the current study includes the determination of antiangiogenic and antioxidant effects of olive phenolic compounds contained in their stable and natural matrix.

Materials and methods

Extract preparation

Olive leaves and fruits were mixed to obtain the aqueous olive tree extract (OTE) by solid-liquid extraction 100% natural, 100% mechanical using water (no use of alcohol, no use of ethanol, no use of chemicals). Actually, OTE is marketed in the world (France, Switzerland, Belgium, ...) under the brandname OLIVIE FORCE / OLIVIE RICHE (see more in www.olivie.ma). The extract consists in a dark brown solution whose major constituents have been determined (table 1). It comes from organic olive trees that are planted in the middle of a rocky desert where trees suffer and trigger a self-defence mechanism where they produce abnormally high quantities of anti-oxydants to survive.

Table 1. Total phenols and major constituents of olive tree extract. Mean \pm standard deviation (*corresponds to the initial first version of OLIVIE FORCE / OLIVIE RICHE*).

| Parameter | Average value (g per100g) |
|--------------------------|---------------------------|
| Density | 1.3 |
| Dry matter | 98.0 \pm 7.8 |
| Mineral matter | 12.9 \pm 0.7 |
| Fat | < 1 |
| Total polyphenols | 7.90 \pm 1.9 |
| Hydroxytyrosol | 2.09 \pm 0.14 |
| ORAC (μ mol Te kg)* | 3848100 \pm 38481 |

*Antioxidant activity determined by ORAC assay (Oxygen Radical Absorbance Capacity).

Cell culture

The HSkMEC (human skin immortalized endothelial cells) was kindly provided by C Kieda (UPR4301 CNRS, Orleans, France) (Kieda 2002), the KB cell line (human nasopharyngeal epidermis carcinoma) was obtained from NCI and HL60 (promyelocytic leukemia) and HT-29 cells (human colon adenocarcinoma) were purchased from ATCC. KB cells in D-MEM, HL60 and HT29 in RPMI and HSkMEC in OPTI-MEM were cultured in complete medium containing 10% (except HSkMEC 2%) fetal bovine serum, penicillin, streptomycin and fungizone in a humidified incubator under 5% CO₂ atmosphere at 37°C. Phenolic compounds (used as reference molecules in this study): oleuropein, hydroxytyrosol, tyrosol, caffeic acid, apigenin, and luteolin were purchased from Sigma Alderich (Paris, France).

MTS assay

HSkMEC, KB and HL60 cell lines were used to determine the cytotoxicity of OTE and reference molecules (phenolic compounds used as standards). Cells were plated in 96-well tissue culture microplates (10^3 to 2.10^3 cells/well in 200 µl of culture medium). After 24 hours of incubation at 37°C under 5% CO₂ atmosphere, cells were treated with increasing concentrations of OTE (diluted in H₂O) and phenolic compounds dissolved in DMSO (1, 10, and 100 µM). Control cells were treated with either water or DMSO (1% of final volume). These operations were carried out using an automated workstation (Biomek 3000 Laboratory Automation Workstation, Beckman-Coulter). After 72h of exposure to olive tree extract or chemicals, MTS reagent (CellTiter 96[®] AQueous One, Promega) was added and incubated for 3 h at 37 °C. Experiments were performed in triplicate: the absorbance was monitored at 490 nm and results were expressed as the inhibition of cell proliferation calculated as the ratio $[(1-(OD_{490} \text{ treated}/OD_{490} \text{ control})) \times 100]$.

FACS analysis

Apoptotic and necrotic cells were analyzed by Fluorescence-Activated Cell Sorting (FACS). KB cells (10^4 cells in 100 µL medium/well in 96-well microplate) were exposed for 24 and 48 hours at 37°C under 5% CO₂ to olive tree extract at different concentrations. 50nM doxorubicin was used as positive control. Microplates were centrifuged to collect floating cells. Adherent cells were trypsinized, washed with PBS, mixed with floating cells and spun down. Cells were resuspended into 100µl

medium prior the addition of 50µl of freshly prepared reaction mixture consisting in the fluorescent agent 7-aminoactinomycin D (7-AAD) (6.5 µL of 1 mg/ml ethanol solution) and recombinant human protein annexin-V-PE (6.5 µL, Bender). After 20 min of incubation in the dark at room temperature, cells were analyzed with a Guava EasyCyte flow cytometer (Millipore). Cells were classified according to their fluorescence and the results were expressed as percentage of cells in each group.

LDH release assay.

Necrosis was estimated through the release of LDH in the culture medium. 20,000 KB cells were incubated for 24 and 48 h in the presence of OTE in 96-well microplates containing 100 µL medium. After centrifugation at 300g for 1 min, 25 µL of culture medium was added with 25 µL Cytotox-ONE reagent (Promega) and kept in the dark at room temperature for 20 min. Fluorescence was recorded (exc 560 nm, em 590 nm): results are expressed as the residual activity in the presence of OTE compared to activity in the presence of vehicle alone.

Caspase activity assay

Caspase activity was determined in KB cells after 24 and 48h of treatment by olive tree extract. Cells ($20 \cdot 10^3$ cells in 180 µl of complete medium per well) were plated in a 96-well microplate under 5% CO₂ atmosphere at 37°C. Lysis buffer (20 µL of a 10x stock solution consisting of 250 mM HEPES buffer, pH 7.5, 5 mM EDTA, 0.5% NP40, 0.1% SDS and 50 mM dithiothreitol) was added before adding caspase-3 substrate (DEVD-AMC at final concentration of 50 µM). The plates were incubated at 37°C and fluorescence was recorded (λ_{ex} 360 nm, λ_{em} 465 nm) after 0, 30, 60, 120, and 180 min. The reaction rate was calculated from the slope of the linear regression (fluorescence vs time) and expressed as the ratio of activation compared to the control (cells treated with DMSO). Doxorubicin (1µM) was used as positive control.

ROS production

HT29 cells were growth in 200 µl RPMI complete medium ($20 \cdot 10^3$ cells/well in black 96-well microplates). After 24 hours, cells were incubated with increasing concentrations of olive tree extract (1 to 100% diluted in H₂O) and 100 µM of reference phenolic compounds (hydroxytyrosol, tyrosol, oleuropein, apigenin, luteolin, caffeic acid dissolved in DMSO). Control cells received DMSO (1% of final

volume). The culture medium was removed and cells were washed with PBS buffer before addition of 100 μ l of RPMI containing 50 mM of 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA). Cells were subsequently incubated for 30 minutes at 37°C before washing with PBS. The fluorescence was monitored after the addition of 100 μ l of PBS (λ_{ex} 485 nm, λ_{em} 530 nm). The positive control was untreated HT29 cells incubated in PBS containing 1 mM of H₂O₂.

In vitro endothelial cell tube formation

In this study, the *in vitro* test of tube formation on Matrigel was used to assess the anti-angiogenic activity of olive tree extract. The collagen matrix (Matrigel) was diluted in ice-cold Opti-MEM without fetal calf serum and 30 μ l of this mix were transferred into a 96-well microplate. The plate was incubated at 37°C under 5% CO₂ atmosphere for 60 min to form the gel. HSkMEC cells (50.10³ cells/well) were seeded in 200 μ l medium containing increasing concentrations of olive tree extract and phenolic compounds (100 μ M). Endothelial cells cultured on Matrigel formed a three-dimensional micro-network of tubes within 16 h of culture. Cell organization was documented photographically with an inverted microscope at \times 10 magnification.

Quantification of pro-angiogenic factors by qRT-PCR

The quantitative RT-PCR was used to measure expression levels of genes coding for the pro-angiogenic factors, vascular endothelial growth factor (VEGF: A, B and C isoforms), hypoxia-inducible factor (HIF-1 α), angiopoietin (ANGPT 1 and 2 isoforms) and VEGF receptor (VEGFR-2) as previously reported (Kieda et al, 2006, Kieda et al, 2013). The cells were lysed and total RNA was extracted from each well of the 96-well microplate using RNA extraction kit according to manufacturer's instructions (Promega, Madison, WI). Two micrograms of RNA were used to prepare cDNA with oligo (dT) and primers of Myeloblastoma Avian Virus (AMV) and reverse transcriptase (Promega). Quantitative real-time PCR was performed using the Roche Light Cycler system and kit FastStart DNA Master SYBR Green-I. Primers were supplied by Qiagen. The data were analyzed with LightCycler 480 Software and values were normalized to the relative amounts of the cDNA of GAPDH gene.

Statistical analysis

All cellular tests were performed in triplicate. Statistical analyzes were performed using GraphPad Prism version 5.00 (GraphPad Inc., San Diego, California). Data were analyzed by analysis of variance (ANOVA Analysis of Variance) followed by post-hoc Dunnet test for normal distributed data or by the Kruskal-Wallis if the sample distribution does not follow the normal law. Differences between groups were considered as statistically significant at $p < 0.05$.

Result

OTE cytotoxicity

A cytotoxic extract is defined as a substance (or mixture of several phytochemical compounds) that is toxic for the cell, acting at several levels from simple metabolic disorders to apoptosis and/or necrosis induction. Actually, studying of plant extract's cytotoxicity is essential to determine appropriate dose prior to its biological uses. Nevertheless, the cytotoxic character is often sought in pharmaceutical field during the research for new drugs with various biological properties. The results presented in this section are those obtained with the commercial MTS test, e.g. the formazan formation followed at 490 nm. The measured absorbance values are directly proportional to the number of living cells and the results were expressed as the inhibition of cell proliferation and viability (Figure 1). Due to its intrinsic color which can quench or conceal measurements, serial dilutions of OTE were applied to cells: the highest concentration 100 indicates that the pure OTE was used at a final concentration of 1 μ l OTE into 100 μ l of medium, whereas the lowest concentration 1 indicates that the pure OTE was diluted 1 to 100 and used at a final concentration of 1 μ l diluted OTE into 100 μ l of medium.




Figure 1. Inhibition of cell viability (%), illustrating the cytotoxicity of olive tree extract in KB cell line.

After 72 hours of treatment, the olive tree extract showed a strong cytotoxic effect on both KB and HSkMEC cell lines. Thus, the olive tree extract was toxic up to 5% dilution with a percentage cell viability inhibition over than 80%. The EC_{50} , concentration of the olive tree extract corresponding to 50% of cell viability inhibition, was subsequently evaluated. KB cells were more sensitive to the olive tree extract ($EC_{50} = 39\%$) than endothelial cells ($EC_{50} = 43\%$), although the observed difference was not significant. Such toxic effect was also observed against other human cancer cells (HL60, MCF-7, HCT116) and non-cancerous cell lines (MRC5 and EPC data not shown). This cytotoxicity activity could be attributed to the phenolic fraction which is present in olive tree extracts (fruits and leaves). Actually, the profile of the olive tree extract revealed a complex mixture of phenolic compounds. Among of them, the components identified as oleuropein, hydroxytyrosol (3',4'-DHPEA) and tyrosol (*p*-DHPEA) were the most represented and flavonoids such as rutin, quercetin, luteolin and apigenin were present in lower concentrations (Laaboudi et al., 2015). For this reason, oleuropein, hydroxytyrosol, tyrosol, apigenin, luteolin, and caffeic acid were used as pure individual molecules for comparison in the current study (Table 2).

Table 2. Inhibition of cell proliferation induced by pure molecules contained in OTE on KB cells. Results are the mean \pm SD of experiments performed in triplicate and expressed as a percentage of inhibition compared to cells treated with DMSO only.

| Compound | Concentration | | |
|----------------|---------------|------------|------------|
| | 100 μ M | 10 μ M | 1 μ M |
| Caffeic acid | 0 \pm 10 | 3 \pm 5 | 6 \pm 6 |
| Apigenin | 83 \pm 1 | 30 \pm 5 | 5 \pm 1 |
| Luteolin | 81 \pm 1 | 22 \pm 6 | 4 \pm 2 |
| Oleuropein | 0 \pm 9 | 0 \pm 1 | 0 \pm 10 |
| Tyrosol | 0 \pm 6 | 5 \pm 3 | 0 \pm 8 |
| Hydroxytyrosol | 11 \pm 11 | 1 \pm 10 | 0 \pm 6 |

Results of table 2 show, surprisingly, that only flavonoids (apigenin and luteolin) were potent inhibitors of KB cell proliferation, while other molecules (tyrosol, hydroxytyrosol, oleuropein and caffeic acid) had no or negligible effect. However flavonoids have no more antiproliferative activity at 1 μ M. Similar results were obtained with other studied cell lines, including HSkMEK (data not shown). This does not exclude the effect on cell proliferation of non-identified compounds (at this stage) present in OTE, including secoiridoids derivatives of hydroxytyrosol and tyrosol especially oleacein (3',4'-DHPEA-EDA) and oleocanthal (*p*-DHPEA-EDA).

FACS analysis of cell death

Apoptosis and necrosis are two distinct and redundant forms of cell death occurring in response to chemical aggression. These two cellular processes featured different and complementary modes of action: apoptosis (or programmed cell death) is a tightly regulated cell death program that plays a pivotal role in a variety of biological processes, generally executed in animal cells via the activation of cellular proteases leading to the cleavage of chromatin into nucleosomal fragments [43], while necrosis involves the destruction of the plasma membrane leading to the release of cytosolic enzymes and cofactors into the external medium. Flow cytometry was used at first to investigate the cell death process: it enabled to discriminate between intact "healthy" cells, apoptotic cells and necrotic cells in an advanced state of degradation.

In line with cytotoxicity experiments, KB cells were treated by diluted OTE (50-1%) for 24 and 48h and analyzed by FACS. According to the fluorescence monitored in FACS, cells can be classified as intact, in early apoptosis, in late apoptosis or necrotic (dead cells). The percentage of cells into each group is shown in Table 3. As it can be inferred from results of table 3, significant changes were observed in cell population distribution after 24 h of exposure to OTE. Intact cell population progressively decreased when the OTE concentration increased, associated with a transient raise in early apoptotic cells at low OTE concentration and a massive outburst of necrotic cells at higher OTE concentrations. After 48h of exposure, the necrotic cell population predominated even with low OTE concentrations to the detriment of intact and early apoptotic cell populations. Therefore it can be concluded that the effects of olive tree extract are dose- and time-dependent.

Table 3. Quantification by FACS of cell death induced in KB cell lines by olive tree extract.

| Cell population (%) | | | | | | | | |
|---------------------|--------|-----|-----------------|-----|----------------|-----|----------|-----|
| Concentration (%) | Intact | | Early apoptotic | | Late apoptotic | | Necrosis | |
| | 24h | 48h | 24h | 48h | 24h | 48h | 24h | 48h |
| DMSO | 90 | 90 | 3.1 | 2.1 | 0.5 | 1.2 | 6.5 | 7.2 |
| Doxo 100 mM | 53 | 32 | 36 | 45 | 0.2 | 1.0 | 11 | 21 |
| OTE 50 | 3.4 | 12 | 4.1 | 1.8 | 0.1 | 0.3 | 92 | 86 |
| OTE 25 | 3.6 | 3.9 | 6.6 | 2.2 | 0.8 | 0.3 | 89 | 94 |
| OTE 10 | 20 | 2.3 | 34 | 8.0 | 0.4 | 0.4 | 46 | 89 |
| OTE 7.5 | 37 | 3.4 | 57 | 12 | 0.2 | 0.4 | 6.0 | 84 |
| OTE 5 | 54 | 19 | 42 | 33 | 0.2 | 0.4 | 3.4 | 47 |
| OTE 2.5 | 86 | 84 | 7.6 | 8.1 | 0.7 | 2.0 | 5.4 | 57 |
| OTE 1 | 83 | 82 | 11 | 9.5 | 0.3 | 0.9 | 4.7 | 7.2 |

Caspases (cysteine-aspartic acid proteases) are regarded as the most common apoptotic effector activated during apoptosis in many cell types. It was reported that caspase-3 is the terminal effector in the apoptotic cascade and is the major contributor to cellular DNA fragmentation.^[33] To go further in the elucidation of the cell death pathway induced by OTE, the catalytic activity of caspase-3 was explored in KB cells treated for 24 and 48 h with OTE at the same concentrations as in FACS analysis. Caspase-3 activity can be determined by the cleavage of the pro fluorescent DEVD peptide which is catalyzed by both caspase-3 and 7, albeit more efficiently by the former. As expected, 1 μ M doxorubicin elicited a potent activation of caspase-3 activity after a 48 h treatment (figure 3A). Although FACS analysis showed the apoptotic effect of olive tree extract at low concentrations (7.5 and 5%), results of figure 3A show that caspase-3 activity was moderately increased after a 48h exposure of KB cells to OTE. This suggest that apoptotic activity of olive tree extract was marginal compared to its necrotic effect and led us to evaluate the release of cytosolic release of lactic dehydrogenase used as a hallmark for necrosis.




Figure 2. Activation of caspase-3 (A) and release of LDH (B) from KB cells after 48h of treatment with olive tree extract, vehicle only (cells) and doxorubicin (1 μ M).

Results are expressed as the relative activity in duplicate experiments \pm SD, compared to activity in untreated KB cells.

As shown in figure 2B, the release of LDH was enhanced after 48h of exposure to increasing OTE concentrations, before to decline at high concentrations, indicative of a complete cell destruction. This confirms that the olive tree extract induces a transient and moderate activation of apoptosis preceding a massive necrotic cell death.

ROS production

The production of reactive oxygen species can be estimated with the DCFH-DA assay using the fluorogenic probe 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA). Briefly, the non-fluorescent DCFH-DA diffuses spontaneously into cells and is deacetylated by cellular esterase to 2',7'-Dichlorodihydrofluorescein (DCFH), which is rapidly oxidized by ROS to the highly fluorescent 2',7'-dichlorodihydrofluoresceine (DCF). The emitted fluorescence intensity (I_{rel}) is proportional to the levels of ROS produced into cell cytosol.




Figure 3. Intracellular ROS production after incubation of HT29 cells for 24h with (A) increasing concentrations of olive tree extract and (B) pure phenol compounds (100 μ M). Results are the mean \pm SD of 3 individual assays. *** ($p < 0.001$), ** ($p < 0.01$) * ($p < 0.05$); ns, not significant from fluorescence produced in cells treated with vehicle only. (1), oleuropein; (2), tyrosol; (3), hydroxytyrosol; (4), apigenin; (5), luteolin; (6), protocatechuic acid; (7) caffeic acid.

In the presence of 1mM H₂O₂, the emitted fluorescence intensity was significantly increased (Figure 3). Conversely, a progressive decrease in the quantity of produced ROS was observed in cells treated with increasing concentrations of OTE to reach an almost complete protection against ROS at highest concentrations. On the other hand, no significant effect on ROS production were noticed with 100µM tyrosol, oleuropein and protocatechuic acid, while hydroxytyrosol, apigenin and luteolin significantly reduced the amount of ROS produced during cell incubation. Thus, the anti-radical effect of olive tree extract could be the result of the antioxidant activity (synergistic action) of these phenolic compounds, although the determination of the role of other compounds (not identified at this stage) is required to confirm these observations.

In vitro anti-angiogenic effect on tube formation.

Endothelial cells grown on an extracellular matrix spontaneously form a capillary network in the presence of growth factors present in the fetal calf serum. This *in vitro* test allow to determine the capacity of a molecule or substance to inhibit the formation of capillary structures. It is considered to be close to *in vivo* tests and to require the proliferation, migration and degradation of the basement membrane of endothelial cells. The photographs taken after a 16h of endothelial cells incubation on Matrigel at 37 °C and 5% CO₂ in the presence of increasing concentrations of olive tree extract and phenolic compounds (100 µM) are presented in figure 4.




Figure 4. Representative photographs of the network structure formed by endothelial cells (HskMEC) on Matrigel in the presence of increasing concentrations of olive tree extract and pure phenolic compounds used at 100 μ M. Control cells are incubated without Matrigel.

Photographs of figure 4 show that endothelial cells grew in complete medium containing 2% FCS as a continuous monolayer (figure 4 control cells) whereas they form capillary structures when laid down on Matrigel. This capacity to form capillary structure was severely impaired in the presence of olive tree extract, suggesting its dose-dependent anti-angiogenic effect. The OTE effect has been observed at concentration 2.5% to be complete at concentration 7.5%. Indeed, referring to the antiproliferative capacity of OTE, the anti-angiogenic activity could be due to a direct cytotoxic effect of OTE on HskMEC cells. However, the results of the cytotoxicity indicate that the olive tree extract diluted at concentration 2.5 does not affect the proliferation and viability of HskMEC cells, whereas the anti-angiogenic effect was evident at this concentration. Therefore the anti-angiogenic effect of pure molecules identified as major phenolic compounds present in olive tree extract was investigated in the same conditions (figure 4). Among these pure compounds hydroxytyrosol, luteolin, apigenin and to a lesser extent caffeic acid at a 100 μ M concentration elicited a marked inhibitory effect on capillary network formation of HskMEC cells. On the

other hand, tyrosol, oleuropein and protocatechuic acid moderately affect the organization of endothelial cells. Thus, we may conclude that the synergistic action of these molecules on endothelial cells might account for the OTE anti-angiogenic effect.

Indeed, the ability of endothelial cells to organize themselves into three-dimensional space to form new capillaries proceeded through cell migration, proliferation and vessel stabilization governed by the secretion of chemotactile factors (called pro-angiogenic factors). Several proteins are involved to support angiogenesis phenomena, such as VEGF factors (A, B and C isoforms), angiotensin (1 and 2), HIF and the VEGF receptor-2. The expression rate of genes coding for these pro-angiogenic factors was, therefore, measured using quantitative real time RT-PCR (Figure 5) and summarized in table 4.

Analytical data plotted on figure 5 were consistent with observation of micro-capillary formation on Matrigel. Basically, the expression of pro-angiogenic genes was significantly affected in the presence of hydroxytyrosol which was the molecule having the highest inhibitory effect on microtube formation, and to a lesser extent by caffeic acid which is a weaker microtube disruptor. The two modest inhibitors of microtube formation tyrosol and oleuropein did not affect the expression of VEGF A and B. It is noteworthy that none of these compounds modify the expression of the VEGF receptor-2 which binds VEGF-A. VHL is a tumor suppressor which binds to HIF-1 α to allow its degradation by the proteasome. Its expression is regulated by HIF-1 α through an HRE element present in its promoter: thus HIF could promote its own degradation by the induction of VHL gene expression (Luczak et al 2011, Karhausen et al 2005). This is consistent with the co-regulation observed in HSkMEC cells exposed to phenolic compounds and presented in figure 5 and table 4.




Figure 5. Effect of phenolic compounds in olive tree extract on the expression of pro-angiogenic agents. HIF Hypoxia Inducible Factor-1 α ; VHL protein of Von Hippel-Lindau; VEGF (A, B and C), Vascular Endothelial Growth Factor; ANGPT (1 and 2), Angiopietin; VEGFR-2, cell receptor for VEGF proteins. Values are expressed as mean \pm SD (n = 3). *** (P < 0.001), ** (P < 0.01) * (P < 0.05); ns, not significant difference.

Table 4. Summary of anti-angiogenic effect of the major polyphenols present in olive tree extract. (+), Inhibitory effect of angiogenesis; (-), No effect.

| Molecule (100 μ M) | HIF | VHL | VEGF- A | VEGF- B | VEGF- C | ANGPT- 1 | ANGPT- 2 | VEGFR- 2 |
|---------------------------|-----|-----|------------|------------|------------|-------------|-------------|-------------|
| Hydroxytyrosol | +++ | +++ | + | + | ++ | ++ | - | - |
| Tyrosol | ++ | - | - | - | + | ++ | - | - |
| Oleuropein | ++ | - | - | - | ++ | + | - | - |
| Caffeic acid | +++ | + | + | + | + | ++ | ++ | - |

All together the lower expression of VEGFs, angiopoietin and HIF-1 α can explain the anti-angiogenic of active phenolic compounds present in the olive tree extract, individually and/or synergistically.

Discussion

The olive tree extract tested in this study, displayed potent cytotoxic activity against human cancer cells through the activation of necrotic and/or apoptotic pathways. This is consistent with several previous studies have reported that crude extracts from olive leaves and phenolic compounds demonstrate high cytotoxic activity against various cancer cell lines derived from blood, colon and breast cancers (Fu et al, 2010;. Hashim et al., 2008; Abaza et al, 2007). This cytotoxic activity was, particularly, remarkable in HER-2 positive breast cancer cells. Olive tree extract reduces the overexpression of HER-2 and decreases resistance to trastuzumab, an anti-HER-2 monoclonal antibody used clinically for breast cancer treatment (Menendez et al., 2007). Independently, Taamalli et al. (2012) have shown that the inhibition of breast cancer cells viability was depended on the polyphenol content in an olive leaves extract.

As expected our olive tree extract is rich in hydroxytyrosol and its secoiridoid derivatives (oleacein and oleocanthal), and in flavones such as luteolin and apigenin. Therefore, the individual effect of these compounds to explain the cytotoxicity toward cancer cells of this crude extract deserves more attention in future studies. Interestingly the whole OTE effect might not result from the abundance of a single compound, but rather from the synergistic interaction between different compounds

including those that are not already identified. In this respect, authors have recently pointed out that food polyphenols who exert their pharmacological effects through synergistic interactions involve the implication of multiple targets (Efferth and Koch, 2011; Wagner, 2011).

Thus, the effects of molecules abundant in OTE like hydroxytyrosol ($50 \mu\text{g}\cdot\text{ml}^{-1}$) and oleuropein ($200 \mu\text{g}\cdot\text{ml}^{-1}$) were demonstrated in MCF-7 cells (Han et al., 2009), and confirmed *in vitro* on a variety of human cancer cell lines together with apigenin (Kim et al, 2011; Bulotta et al, 2011; Fabiani et al 2006, 2002), and *in vivo* in tumor-bearing mice (Hamdi and Castellon, 2005). Moreover, many epidemiological studies suggest a potential correlation between the consumption of olive products (olive fruits and olive oil) and a low incidence of breast cancer in Mediterranean area, suggesting a protective effect of olive compounds against cell proliferation e.g. cancer (Corona et al., 2007). Thus, we can conclude that the OTE's cytotoxic effect is mainly due to these phenolic compounds.

ROS play a crucial role in tumorigenesis as suggested by Sabharwal and Schumacker (2014) who reported that signaling pathway(s) triggered by ROS contributes to proliferation and tumor survival of many cancers. These authors propose that improving mitochondria potential redox is a promising target for future cancer therapies. In accordance with Nardi et al. (2014), we reported herein that olive tree extract and some of its phenolic components showed elevated antioxidant activity in HT-29 cells, giving them a central role in the fight against carcinogenic risks. Likewise, Deiana et al. (1999) reported that hydroxytyrosol reduced biochemical effects of peroxynitrite, such as the deamination of adenine and guanine in certain cell lines. ROS generated during oxidative stress may be directly involved in DNA damage and/or in the intrinsic apoptotic pathway induction (mitochondria-dependent). Thus, the intracellular accumulation of ROS may be involved in the partial disruption of the mitochondrial membrane causing the release of mitochondrial material (cytochrome-c, pro-apoptotic factors) in the cytosol, finally leading to the activation of the caspase cascade (Eloy et al., 2012). Moreover, cellular ROS production by different metabolic pathways has been associated with cell aging phenomenon call the "radical" hypothesis of aging (Barouki, 2006). Thus, we can postulate the antioxidant activity of OTE and these phenolic compounds has a double effect, i) the elimination of carcinogenic ROS and ii) the prevention of cell aging.

Angiogenesis is a critical process implicated in the proliferation and dissemination of cancer cells by metastasis. The fast and limitless multiplication of cancer cell requires an important and incremental supply in oxygen and nutrients through the blood circulation. The spreading of circulating cancer cells is highly facilitated by the density of blood vessels and the cell aggressiveness enabling the metastasis formation.

Angiogenesis is under the control of multiple regulation pathways, among them are the vascular endothelial growth factors which stimulate vasculogenesis and angiogenesis (Jiang et al., 2000) and facilitate the formation of new capillary vessels in association with angiopoietins reviewed by Rak et al., 2000. The transcription of VEGF is stimulated by Hypoxia Inducing Factor 1- α present in the cytosol as a heterodimer with VHL allowing its degradation by the proteasome. The key role of VEGFs was also highlighted by a clinical report which positively correlated polymorphism in the VEGF-A gene (-1154 A/G, -2578 C/A and -460 T/C) and the lower susceptibility to develop breast cancer in Moroccan individuals (Rahoui et al., 2014).

Data reported herein clearly demonstrate the inhibitory effect of olive tree extract and of some of its phenolic components (hydroxytyrosol, apigenin, luteolin and caffeic acid) on the formation of new capillaries from human endothelial cells. This inhibition was consistent with previous observations made on various human cells, animal tumors and *in silico* (Ambasta et al., 2015; Simon Silvan et al., 2013; Fortes et al., 2012; Gacche et al., 2011; Engelmann et al. 2002). Two mechanisms could explain the anti-angiogenic effect of olive tree extract and phenolic compounds: a severe restriction of cell proliferation reducing the number of endothelial cells required for building the capillary structure and a lack of pro-angiogenic factors production thus restricting cell migration and assembly. Clearly, hydroxytyrosol oleuropein and caffeic acid had no effect on cell proliferation, but are potent down-regulator of HIF, VEGF and angiopoietin suggesting their direct anti-angiogenic effect. Additionally, Kim et al (2011) had shown that apigenin has an anti-angiogenesis effect through a down-regulation of VEGF and MMP-8 release in hepatocellular carcinoma cells. Therefore, OTE's effect is the result of the cumulative effect of individual phenolic compounds reducing the cell supply and thus restricting the tumor growth.

On the other hand, an excessive angiogenesis ensues from progression of inflammatory process in synovium, and leads to pannus proliferation and rheumatoid

arthritis symptoms complication. Angiogenesis is a major contributor in the development and maintain of inflammation in rheumatoid arthritis (Semerano et al., 2011), and a correlation between rheumatoid arthritis progression and VEGF level has been observed in patient with RA (Sone et al., 2001). Thus, the inhibition of angiogenesis could be a potential therapeutic target for the treatment of rheumatoid arthritis (Ghanam et al., 2015). However, the VEGF-dependent antiangiogenic effect of olive tree polyphenols, in both cancer and rheumatoid arthritis diseases, need to be confirmed in future *in vivo* studies.

Conclusion

The cellular assays have demonstrated the effectiveness of phenolic olive tree extract as an antioxidant in cells exposed to oxidative stress. This extract has also demonstrated a strong VEGF-dependent antiangiogenic activity. Indeed, ROS elimination, apoptosis and/or necrosis activation and angiogenesis inhibition make this extract an excellent agent to prevent cancer diseases. The characterization of bioactive molecules in this extract, as well as the contribution of each to the global biological activity, could provide the chemical basis for the development of new anticancer drugs. Further studies are needed to extent the antiangiogenic activity of phenolic compounds derived from studied olive tree extract (hydroxytyrosol, oleacein and oleocanthal) in case of angiogenesis-dependent diseases.

References

Semerano L, Clavel G, Assier E, Denys A, Boissier MC. Blood vessels, a potential therapeutic target in rheumatoid arthritis? *Jt Bone Spine* 2011; 78: 118-123.

Sone H, Kawakamib Y, Sakauchic M, Nakamurad Y, Takahashia A, Shimano H et al. Neutralization of Vascular Endothelial Growth Factor Prevents Collagen-Induced Arthritis and Ameliorates Established Disease in Mice. *Biochem Biophys Res Commun* 2001; 281: 562–568.

Abaza L, Talorete T.P.N, Yamada P, Kurita Y, Zarrouk M, Isoda H. Tunisian Gerboui olive leaf extract induces growth inhibition and differentiation of human leukemia HL-60 cells. *J Biosci Biotechnol Biochem* (2007); 7: 1306–1312.

Ambasta RK, Jha SK, Kumar D, Sharma R, Jha NK, Kumar P. Comparative study of anti-angiogenic activities of luteolin, lectin and lupeol biomolecules. *J Transl Med.* 2015, 13, 307.

Barouki R. Stress oxydant et vieillissement. *Medecine Sciences* (2006); 22: 266-72.

Benavente-García O, Castillo J, Lorente J, Ortuño A, Del Río J.A. Antioxidant activity of phenolics extracted from *Olea europea* L. leaves. *Food Chem* (2000); 68: 457-462.

Bulotta S, Corradino R, Celano M, D'Agostino M, Maiuolo J, Oliverio M, Procopio A, Iannone M, Rotiroti D, Russo D. Antiproliferative and antioxidant effects on breast cancer cells of oleuropein and its semisynthetic peracetylated derivatives. *Food Chem* (2011); 127: 1609–1614.

Corona G, Tzounis X, Dessi M.A, Deiana M, Debnam E.S, Visioli F, Spencer J.P.E. The fate of olive oil polyphenols in the gastrointestinal tract: implications of gastric and colonic microflora dependent biotransformation. *Free Radical Res* (2006); 40: 647–658.

Covas M. I. Olive oil and the cardiovascular system. *Pharmacol. Res.* 2007, 55, 452 175–186.

Covas MI, de la Torre K, Farre-Albaladejo M, Kaikkonen J, Fito M, Lopez-Sabater C. (2006). Postprandial LDL phenolic content and LDL oxidation are modulated by olive oil phenolic compounds in human. *Free Radic Biol Med*, 40, 608–16.

Deiana M, Aruoma OI, Bianchi MDLP, Spencer JPE, Kaur H, Halliwell B, Aeschbach R, Banni S, Dessi MA, Corongiu FP (1999). Inhibition of peroxynitrite dependent DNA base modification and tyrosine nitration by the extra virgin olive oil-derived antioxidant hydroxytyrosol. *Free Radical Biology and Medicine.* 26: 762–769.

Efferth T, Koch E. Complex interactions between phytochemicals. The multi-target therapeutic concept of phytotherapy. *Curr Drug Targets* (2011); 12: 122–132.

Eloy L, Jarrousse AS, Teyssot ML, Gautier A, Morel L, Jolival C, Cresteil T, Roland S. Anticancer Activity of Silver–N-Heterocyclic Carbene Complexes: Caspase-Independent Induction of Apoptosis via Mitochondrial Apoptosis-Inducing Factor (AIF). *ChemMedChem* 2012, 7, 805 – 814.

Engelmann C, E. Blot, Y. Panis, S. Bauer, V. Trochon, H. J. Nagy, H. Lu, C. Soria. Apigenin – strong cytostatic and anti-angiogenic action in vitro contrasted by lack of efficacy in vivo. *Phytomedicine* 9: 489–495, 2002.

Estruch R, Martinez-Gonzalez MA, Corella D, Salas-Salvadó J, Ruiz-Gutiérrez V, Covas MI. (2006). Effects of a Mediterranean-Style Diet on Cardiovascular Risk Factors: A Randomized Trial. *Ann Intern Med*, 145, 1-11.19.

Evagelia T, Harris N. Lazarides and Konstantinos B (2004). Olive Mill Wastewater Treatment. Petrotos Aristotle University of Thessaloniki, Department of Food Science and Technology, 54006, Thessaloniki, Greece.

Fabiani R, De Bartolomeo A, Rosignoli P, Servili M, Montedoro G.F, Morozzi G. Cancer chemoprevention by hydroxytyrosol isolated from virgin olive oil through G1 cell cycle arrest and apoptosis. *Eur J Cancer Prev* (2002); 11: 351–358.

Fabiani R, De Bartolomeo A, Rosignoli P, Servili M, Montedoro G.F, Morozzi G Virgin olive oil phenols inhibit proliferation of human promyelocytic leukemia cells (HL60) by inducing apoptosis and differentiation. *J Nutr* (2006); 136: 614–619.

Fortes C, García-Vilas J, Quesada A, Medina M. Evaluation of the anti-angiogenic potential of hydroxytyrosol and tyrosol, two bio-active phenolic compounds of extra virgin olive oil, in endothelial cell cultures. *Food Chem* 2012; 134: 134–140.

Fu S, Arráez-Roman D, Segura-Carretero A, Menéndez J.A, Menéndez-Gutiérrez M.P, Micol V, Fernández-Gutiérrez A. Qualitative screening of phenolic compounds in olive leaf extracts by hyphenated liquid chromatography and preliminary evaluation of cytotoxic activity against human breast cancer cells. *Anal Bioanal Chem* (2010); 397: 643–654.

Gacche RN, Shegokar HD, Gond DS, Yang Z, Jadhav AD. Evaluation of selected flavonoids as antiangiogenic, anticancer, and radical scavenging agents: an experimental and in silico analysis. *Cell Biochem Biophys*. 2011, 61, 651-63.

Ghanam J., Laaboudi W., Benlemlih M. Effect of rich polyphenols olive tree extract on inflammation and pain in patients with rheumatoid arthritis: a 8-week randomized, double blind, placebo-controlled clinical trial. *American Journal of Biological and Pharmaceutical Research*. 2015; 2(2): 51-61.

Giacosa A, Barale R, Bavaresco L, Gatenby P, Gerbi V, Janssens J, Johnston B, Kas K, La Vecchia C, Mainguet P, et al. Cancer prevention in Europe: the Mediterranean diet as a protective choice. *Eur J Cancer Prev* 2013; 22:90-5.

Hamdi K.H & Castellon R. Oleuropein, a non-toxic olive iridoid, is an anti-tumor agent and cytoskeleton disruptor. *Biochem Bioph Res Co* (2005); 334: 769-778.

Han J, Talorete TP, Yamada P, Isoda H. Anti-proliferative and apoptotic effects of oleuropein and hydroxytyrosol on human breast cancer MCF-7 cells. *Cytotechnology* (2009); 59: 45–53.

Hashim Y.Z, Rowland I.R, McGlynn H, Servili M, Selvaggini R, Taticchi A, Esposto S, Montedoro G, Kaisalo L, Wähälä K, Gill C.I. Effects of olive oil phenolics on invasion in human colon adenocarcinoma cells *in vitro*. *Int J Cancer* (2008); 122: 495–500.

Japón-Luján R, Luque de Castro M.D. Small branches of olive tree: A source of biophenols complementary to olive leaves. *J Agric Food Chem* (2007); 55: 4584-4588.

Jiang BH, Zheng JZ, Aoki M, Vogt PK. Phosphatidylinositol 3-kinase signaling mediates angiogenesis and expression of vascular endothelial growth factor in endothelial cells. *Proc Natl Acad Sci USA* 97: 1749-1753, 2000.

Karhausen J, Kong T, Narravula S, Colgan SP. Induction of the von Hippel-Lindau tumor suppressor gene by late hypoxia limits HIF-1 expression. *J. Mol. Biochem*, 2005, 95, 1264-1275.

Kieda C, El Hafny-Rahbi B, Collet G, Lamerant-Fayel N, Grillon C, Guichard A, Dulak J, Jozkowicz A, Kotlinowski J, Fylaktakidou KC, Vidal A, Auzeloux P, Miot-Noirault E, Beloeil JC, Lehn JM, Nicolau C. Stable tumor vessel normalization with pO₂ increase and endothelial PTEN activation by inositol tris pyrophosphate brings novel tumor treatment. *J Mol Med (Berl)*. 2013, 91, 883-899.

Kieda C, Greferath R, Crola da Silva C, Fylaktakidou KC, Lehn JM, Nicolau C. Suppression of hypoxia-induced HIF-1 α and of angiogenesis in endothelial cells by myo-inositol tris pyrophosphate-treated erythrocytes. *Proc Natl Acad Sci U S A*. (2006) 103, 15576-81.

Kieda C, Paprocka M, Krawczenko A, Załecki P, Dupuis P, Monsigny M, Radzikowski C, Duś D. New human microvascular endothelial cell lines with specific adhesion molecules phenotypes. *Endothelium*. (2002), 9, 47-61.

Killeen M, Linder M, Pontoniere P, Crea R. NF- κ B signaling and chronic inflammatory diseases: exploring the potential of natural products to drive new therapeutic opportunities. *Drug Discov Today* (2014); 19: 373-378.

Kim B.R, Jeon Y.K, Nam M.J. A mechanism of apigenin-induced apoptosis is potentially related to anti-angiogenesis and anti-migration in human hepatocellular carcinoma cells. *Food Chem Toxicol* (2011); 49: 1626–1632.

LeGendre O, Breslin P A S, Foster D A. (-)-Oleocanthal rapidly and selectively induces cancer cell death via lysosomal membrane permeabilization (LMP). *Molecular & Cellular Oncology*, 2:4, e1006077.

Łuczak MW, Roszak A, Pawlik P, H Kedzia H, M Lianeri M, Jagodzinski PP. Increased expression of HIF-1A and its implication in the hypoxia pathway in primary advanced uterine cervical carcinoma. *Oncol. Rep.*, 2011, 26, 1259-1264.

Mario-Casas E, Covas MI, Farre M, Fito M, Ortuno J, Weinbrenner T, Roset P, de la Torre R. Hydroxytyrosol disposition in humans. *Clinical Chem* (2003a); 49: 945–952.

Menendez J.A, Vazquez-Martin A, Colomer R, Brunet J, Carrasco-Pancorbo A, Garcia-Villalba R, Fernandez-Gutierrez A, Segura-Carretero A. Olive oil's bitter principle reverses acquired autoresistance to trastuzumab (Herceptin) in HER2-overexpressing breast cancer cells. *BMC Cancer* (2007); 7: 80.

Micol V, Caturla N, Pérez-Fons L, Más V, Pérez L, Estepa A. The olive leaf extract exhibits antiviral activity against viral haemorrhagic septicaemia rhabdovirus (VHSV). *Antiviral Res* (2005); 66: 129-136.

Montedoro G.F, Servili M, Baldioli M, Selvaggini R, Miniati E, Maccchioni A. Simple and hydrolyzable compounds in virgin olive oil. 3. Spectroscopic characterizations of the secoiridoid derivatives. *J Agric Food Chem* (1993); 41: 2228–2234.

Nardi M, Bonacci S, De Luca G, Maiuolo J, Oliverio M, Sindona G, Procopio A. Biomimetic synthesis and antioxidant evaluation of 3,4-DHPEA-EDA [2-(3,4-hydroxyphenyl) ethyl (3S,4E)-4-formyl-3-(2-oxoethyl) hex-4-enoate]. *Food Chem*

(2014); 162: 89–93.

Raederstorff D, Wang-schmidt Y, Wertz K (2010). Use of hydroxytyrosol as anti-aging agent. Pub. No. : US 2010 0130621 A1.

Rahoui J, Laraqui A, Sbitti Y, Touil N, Ibrahim A, Ghrab B, Al Bouzidi A, Moussaoui D, Rahali, Dehayni M, Ichou M, Zaoui F, Mrani S. Investigating the association of vascular endothelial growth factor polymorphisms with breast cancer: a Moroccan case–control study. *Medical Oncology* 2014, 31:193.

Rak J, Yu JL, Klement G, Kerbel RS. Oncogenes and angiogenesis: signaling three-dimensional tumor growth. *Journal of Investigative Dermatology Symposium Proceedings* (2000) 5, 24–33.

Rashmi K. Ambasta, Saurabh Kumar Jha, Dhiraj Kumar, Renu Sharma, Niraj Kumar Jha, Pravir Kumar. Comparative study of anti-angiogenic activities of luteolin, lectin and lupeol biomolecules. *J Transl Med* (2015) 13:307.

Richard N, Arnold S, Hoeller U, Kilpert C, Wertz K, Schwager J. Hydroxytyrosol Is the Major Anti-Inflammatory Compound in Aqueous Olive tree extracts and Impairs Cytokine and Chemokine Production in Macrophages. *Planta Medica* (2011); 77: 1890-1897.

Sabharwal S. & Schumacker P.T. Mitochondrial ROS in cancer: initiators, amplifiers or an Achilles' heel? *Nat Rev Cancer* (2014); 14: 709–721.

Scoditti E, Calabriso N, Massaro M, Pellegrino M, Storelli C, Martines G. (2012). Mediterranean diet polyphenols reduce inflammatory angiogenesis through MMP -9 and COX-2 inhibition in human vascular endothelial cells: a potentially protective mechanism in atherosclerotic vascular disease and cancer. *Arch Biochem Biophys*, 527, 81–89.

Silvan S, Manoharan S. Apigenin prevents deregulation in the expression pattern of cell-proliferative, apoptotic, inflammatory and angiogenic markers during 7,12-dimethylbenz[a]anthracene-induced hamster buccal pouch carcinogenesis. *Arch Oral Biol*. 2013, 58, 94-101.

Taamalli A, Arráez-Román D, Barrajón-Catalán E, Ruiz-Torres V, Pérez-Sánchez A, Herrero M, Ibáñez E, Micol V, Zarrouk M, Segura-Carretero A, Fernández-Gutiérrez

A. Use of advanced techniques for the extraction of phenolic compounds from Tunisian olive leaves: Phenolic composition and cytotoxicity against human breast cancer cells. *Food Chem Toxicol* (2012); 50: 1817–1825.

Visioli F, Galli C, Bornet F, Mattei A, Patelli R, Galli G, et al. Olive oil phenolics are dose-dependently absorbed in humans. *FEBS Lett* (2000); 468:159–160.

Wafa Laaboudi, Jamal Ghanam, Mohammed Merzouki and Mohamed Benlemlih. Eco-Extraction of Phenolic Compounds from Moroccan Olive Fruits and Leaves and their Potential use as Antimicrobial Agents. *European Journal of Scientific Research*. (2015); 132: 255-265.

Wagner H. Synergy research: approaching a new generation of phytopharmaceuticals. *Fitoterapia* (2011); 82: 34–37.